

# The COLchicine Cardiovascular Outcomes Trial

**Jean-Claude Tardif MD**, Simon Kouz, MD, David D. Waters MD,  
Olivier F. Bertrand MD PhD, Rafael Diaz MD, Aldo P. Maggioni MD,  
Fausto J. Pinto MD PhD, Reda Ibrahim MD, Habib Gamra MD,  
Ghassan S. Kiwan MD, Colin Berry MD PhD, Jose Lopez-Sendon MD,  
Petr Ostadal MD PhD, Wolfgang Koenig MD, Denis Angoulvant MD,  
Jean C. Grégoire MD, Marc-André Lavoie MD, Marie-Pierre Dubé PhD,  
David Rhainds PhD, Mylène Provencher PhD, Lucie Blondeau MSc,  
Andreas Orfanos MBBCh, Philippe L. L'Allier MD,  
Marie-Claude Guertin PhD, François Roubille MD, PhD

on behalf of the COLCOT Investigators

Montreal Health Innovations  
Coordinating Center

A Division of the Montreal Heart Institute



MONTREAL  
**HEART**  
INSTITUTE

# Background



- **Experimental and clinical evidence support the role of inflammation in atherosclerosis and its complications.**
- **The search for a widely used anti-inflammatory treatment that may reduce the risk of atherosclerotic events in patients with coronary artery disease continues.**
- **Colchicine is an orally administered, potent anti-inflammatory medication currently indicated for gout and pericarditis.**
- **COLCOT was conducted in patients with a recent myocardial infarction to evaluate the effects of colchicine on cardiovascular outcomes and its long-term safety and tolerability.**

# Study design



Post-myocardial infarction  $\leq 30$  days (n=4745 patients)  
On statin, anti-platelet agents,  $\pm$ RAASi,  $\pm$ BB

Treated according to national guidelines  
PCI completed if planned

Colchicine 0.5 mg  
daily \*

Placebo  
daily \*

Primary composite endpoint: Time to first of CV death, cardiac arrest, MI, stroke, or urgent hospitalization for angina requiring coronary revascularization

Secondary endpoints: Components of primary;  
composite of CV death, cardiac arrest, MI or stroke; total mortality

# Patient characteristics


	<b>Colchicine (N=2366)</b>	<b>Placebo (N=2379)</b>
Age - years	60.6±10.7	60.5±10.6
Female sex - no. (%)	472 (19.9%)	437 (18.4%)
Caucasian - no. (%)	1350/1850 (73.0%)	1329/1844 (72.1%)
Body-mass index - kg/m <sup>2</sup>	28.2±4.8	28.4±4.7
Smoking - no. (%)	708 (29.9%)	708 (29.8%)
Hypertension - no. (%)	1185 (50.1%)	1236 (52.0%)
Diabetes - no. (%)	462 (19.5%)	497 (20.9%)
Prior MI - no. (%)	370 (15.6%)	397 (16.7%)
Prior PCI - no. (%)	392 (16.6%)	406 (17.1%)
Prior CABG - no. (%)	69 (2.9%)	81 (3.4%)
Prior heart failure - no. (%)	48 (2.0%)	42 (1.8%)
Prior stroke/TIA - no. (%)	55 (2.3%)	67 (2.8%)

# Patient characteristics



	<b>Colchicine (N=2366)</b>	<b>Placebo (N=2379)</b>
Index MI to randomization - days	13.4 ± 10.2	13.5 ± 10.1
PCI for index MI - no. (%)	2192/2364 (92.7%)	2216/2375 (93.3%)
Aspirin - no. (%)	2334 (98.6%)	2352 (98.9%)
Other anti-platelet agent - no. (%)	2310 (97.6%)	2337 (98.2%)
Statin - no. (%)	2339 (98.9%)	2357 (99.1%)
Beta-blocker - no. (%)	2116 (89.4%)	2101 (88.3%)


# Primary efficacy endpoint



## No. at Risk

Colchicine	2366	2284	1868	1230	628	153	0
Placebo	2379	2261	1854	1224	622	144	0

# Primary efficacy endpoint




## No. at Risk

Colchicine	2366	2284	1868	1230	628	153	0
Placebo	2379	2261	1854	1224	622	144	0

# Major Clinical Outcomes

Clinical Outcome	Colchicine	Placebo	Hazard Ratio	P
Intent-to-treat population	N=2366	N=2379	(95% CI)	Value
<u>Primary composite endpoint</u> - no. (%)	<u>131 (5.5%)</u>	<u>170 (7.1%)</u>	<u>0.77 (0.61-0.96)</u>	<u>0.02</u>
CV death - no. (%)	20 (0.8%)	24 (1.0%)	0.84 (0.46-1.52)	
Resuscitated cardiac arrest - no. (%)	5 (0.2%)	6 (0.3%)	0.83 (0.25-2.73)	
Myocardial infarction - no. (%)	89 (3.8%)	98 (4.1%)	0.91 (0.68-1.21)	
Stroke - no. (%)	5 (0.2%)	19 (0.8%)	0.26 (0.10-0.70)	
Urgent hospitalization for angina requiring revascularization - no. (%)	25 (1.1%)	50 (2.1%)	0.50 (0.31-0.81)	
<u>Secondary composite endpoint</u> - no. (%)	111 (4.7%)	130 (5.5%)	0.85 (0.66-1.10)	
Death - no. (%)	43 (1.8%)	44 (1.8%)	0.98 (0.64-1.49)	
DVT or pulmonary embolus - no. (%)	10 (0.4%)	7 (0.3%)	1.43 (0.54-3.75)	
Atrial fibrillation - no. (%)	36 (1.5%)	40 (1.7%)	0.93 (0.59-1.46)	

# Primary efficacy endpoint



**No. at Risk**

Colchicine	2260	2197	1791	1169	601	140	0
Placebo	2270	2169	1778	1173	596	135	0

# Major Clinical Outcomes

Clinical Outcome	Colchicine	Placebo	Hazard Ratio	P
Per-protocol population	N=2260	N=2270	(95% CI)	Value
<u>Primary composite endpoint</u> - no. (%)	115 (5.1%)	162 (7.1%)	0.71 (0.56-0.90)	<b><u>0.004</u></b>
CV death - no. (%)	19 (0.8%)	23 (1.0%)	0.83 (0.45-1.53)	
Resuscitated cardiac arrest - no. (%)	5 (0.2%)	5 (0.2%)	1.00 (0.29-3.46)	
Myocardial infarction - no. (%)	77 (3.4%)	92 (4.1%)	0.84 (0.62-1.14)	
Stroke - no. (%)	5 (0.2%)	19 (0.8%)	0.26 (0.10-0.71)	
Urgent hospitalization for angina requiring revascularization - no. (%)	22 (1.0%)	47 (2.1%)	0.47 (0.28-0.78)	

# Total (First + Recurrent) Primary Endpoint Events (ITT)

Endpoint / Model		Colchicine N=2366	Placebo N=2379	Hazard / Rate Ratio (95% CI)
Total number of primary endpoint events		154	223	
Rate of primary endpoint events per 100 patient-months		0.29	0.42	
Negative binomial model				0.66 (0.51; 0.86)
Andersen-Gill model				0.69 (0.54; 0.88)
Wei-Lin-Wessfeld model	1 <sup>st</sup> event			0.77 (0.61; 0.96)
Wei-Lin-Wessfeld model	2 <sup>nd</sup> event			0.73 (0.48; 1.11)
Wei-Lin-Wessfeld model	3 <sup>rd</sup> event			0.64 (0.37; 1.10)
Wei-Lin-Wessfeld model	Average			0.77 (0.61; 0.96)

# Adverse events



Safety population	Colchicine (N=2330)	Placebo (N=2346)	P Value
Any related AE - no. (%)	372 (16.0%)	371 (15.8%)	0.89
Any SAE - no. (%)	383 (16.4%)	404 (17.2%)	0.47
Gastro-intestinal AE - no. (%)	408 (17.5%)	414 (17.6%)	0.90
Gastro-intestinal SAE – no. (%)	46 (2.0%)	36 (1.5%)	0.25
Diarrhea AE - no. (%)	225 (9.7%)	208 (8.9%)	0.35
Nausea AE - no. (%)	43 (1.8%)	24 (1.0%)	<b>0.02</b>
Flatulence AE - no. (%)	15 (0.6%)	5 (0.2%)	<b>0.02</b>
GI haemorrhage AE - no. (%)	7 (0.3%)	5 (0.2%)	0.56
Infection SAE - no. (%)	51 (2.2%)	38 (1.6%)	0.15
Pneumonia SAE - no. (%)	21 (0.9%)	9 (0.4%)	<b>0.03</b>
Septic shock SAE - no. (%)	2 (0.1%)	2 (0.1%)	0.99
HF hospitalization - no. (%)	25 (1.1%)	17 (0.7%)	0.21
Cancer - no. (%)	43 (1.8%)	46 (2.0%)	0.77
Anemia - no. (%)	14 (0.6%)	10 (0.4%)	0.40
Leukopenia - no. (%)	2 (0.1%)	3 (0.1%)	0.66
Thrombocytopenia - no. (%)	3 (0.1%)	7 (0.3%)	0.21


# Limitations

- **The duration of follow-up was relatively short at approximately 23 months. The risks and benefits of longer-term treatment with colchicine were not evaluated.**
- **Although the inclusion of 4745 patients was sufficient to demonstrate a significant benefit on the primary composite efficacy endpoint, a larger trial could have allowed a better assessment of individual endpoints and subgroups and the risks associated with colchicine.**

## Conclusion

- **Colchicine 0.5 mg/day significantly reduces the risk of first and total ischemic cardiovascular events by 23% and 34% respectively compared to placebo in patients with a recent myocardial infarction.**
- **Rates of adverse effects were low, including a small increase in pneumonias (0.9 vs. 0.4%) but no significant increase in diarrhea with colchicine, on background therapy with aspirin, a 2nd antiplatelet agent and a statin in 99, 98 and 99% of patients.**
- **The COLCOT results apply to patients who have recently suffered a myocardial infarction. Further research is needed to assess the benefits of colchicine in other high-risk patients.**

# COLCOT-T2D – Study design





The NEW ENGLAND  
JOURNAL of MEDICINE



ORIGINAL ARTICLE

## Efficacy and Safety of Low-Dose Colchicine after Myocardial Infarction

Jean-Claude Tardif, M.D., Simon Kouz, M.D., David D. Waters, M.D.,  
Olivier F. Bertrand, M.D., Ph.D., Rafael Diaz, M.D., Aldo P. Maggioni, M.D.,  
Fausto J. Pinto, M.D., Ph.D., Reda Ibrahim, M.D., Habib Gamra, M.D.,  
Ghassan S. Kiwan, M.D., Colin Berry, M.D., Ph.D., José López-Séndon, M.D.,  
Petr Ostadal, M.D., Ph.D., Wolfgang Koenig, M.D., Denis Angoulvant, M.D.,  
Jean C. Grégoire, M.D., Marc-André Lavoie, M.D., Marie-Pierre Dubé, Ph.D.,  
David Rhoads, Ph.D., Mylène Provencher, Ph.D., Lucie Blondeau, M.Sc.,  
Andreas Orfanos, M.B., B.Ch., Philippe L. L'Allier, M.D.,  
Marie-Claude Guertin, Ph.D., and François Roubille, M.D., Ph.D.

Montreal Health Innovations  
*Coordinating Center*

A Division of the Montreal Heart Institute



MONTREAL  
**HEART**  
INSTITUTE